

## Arkansas Department of Health

4815 West Markham Street • Little Rock, Arkansas 72205-3867 • Telephone (501) 661-2000 Governor Asa Hutchinson Nathaniel Smith, MD, MPH, Secretary of Health

March 17, 2020

## Clinical Guidance for Management of Patients with Confirmed COVID-19

Interim clinical guidance for the management of patients with confirmed COVID-19 (based on current information) is available at the CDC and WHO websites:

https://www.cdc.gov/coronavirus/2019-ncov/hcp/clinical-guidance-management-patients.html

https://www.who.int/publications-detail/clinical-management-of-severe-acute-respiratory-infection-when-novel-coronavirus-(ncov)-infection-is-suspected

We urge you to familiarize yourself with this information as you will likely care for a confirmed COVID-19 patient. A few highlights are included below:

- Incubation Period of SARS-CoV-2: median range of 4 days (IQR of 2 to 7 days) with a maximum of 14 days.
- Most common symptoms at presentation: fever, cough, myalgia, fatigue, shortness of breath.
- Risk factors for severe illness include older patients (usually 65 or greater) and those with chronic medical conditions such as cardiovascular disease, diabetes, chronic respiratory disease, hypertension, and malignancy.
- Illness can range from asymptomatic to mild URI symptoms to severe illness requiring mechanical ventilation, and sometimes death.
- Mild disease can sometimes be managed at home but this is at the discretion of the healthcare personnel evaluating patient.
  - o Disease may worsen in the second week after infection.
  - Older patients and those with chronic medical conditions, as stated above, are at higher risk for progression of disease.
- Corticosteroids should be avoided as they can prolong viral replication.

Finally, we would also like to reiterate the importance of rapid implementation of your facility's infection control and prevention measures.

Updates will be provided as more information and guidance becomes available.